

Compensatory measures for wildlife conservation: testing the effect of deadwood and cavity supply on cavity users in managed boreal forest

Samuel Dufour-Pelletier¹, Junior A. Tremblay², Jacques Ibarzabal¹

¹Université du Québec à Chicoutimi ²Environment and Climate Change Canada

UQAC
Université du Québec
à Chicoutimi



Forêts, Faune
et Parcs
Québec



Environment and
Climate Change Canada



CONTEXT

Forest management leads to a decrease in the proportion of **old forest**^{1,2,3} and in the quality/amount of **deadwood**^{4,5,6}. The simplification of the internal forest structure and the overabundance of second growth forest stands may be a limiting factor for associated wildlife^{7,8,9}.





Figure 1. Managed forest in study area

Ecosystem-based forest management is the established approach under Quebec's new forest regime to address environmental challenges.

Anthropogenic supply of standing deadwood^{10,11} and cavities^{12,13,14} may have a positive impact on associated biodiversity.


Objective

Determine whether an **anthropogenic compensatory measure** (deadwood and cavity supply) in a managed boreal forest can emulate attributes of an old-growth forest and allow the presence and reproduction of deadwood associated species.




STUDY AREA

- Eastern Canadian boreal forest under management
- Before-After– Control-Impact (BACI)¹⁵ (2015-2016)
- 5 experimental blocks of 6 sampling units (SU)
- 1 treatment per SU per block
 - Deadwood distributed uniformly
 - Deadwood distributed in patch
 - Deadwood distributed uniformly with cavities
 - Deadwood distributed in patch with cavities
 - Cavities
 - Control
- Forest stands
 - Black spruce
 - 50 - 70 years old
 - ≥ 4 hectares
 - Similar tree composition
 - Distanced >1.5 km
 - Crossed by a forest road
 - Excluding perturbations and watercourse




METHODS

- Bird surveys (Springs 2015 and 2016)
 - Point counts (15 min.)
 - Playback: BOCH, RBNU, ATTW, BBWO, NOFL, BOOW, NHOW
 - Recordings: omnidirectional microphone/ TASCAM recorder
 - Unlimited radius
 - Post sampling identification
- Saproxylic insects surveys (Springs 2015 and 2016)
 - Trunk Windows Trap
 - Ethanol [70%]
- Nest-boxes survey during breeding season (Spring 2016)
 - Breeding success
 - Motion-detection cameras for occupied nest-boxes



- Vegetation plots
 - Trees (400m²)
 - Natural deadwood (1250m²)
- Woodpecker feeding marks
 - Number of holes
 - Scaling surface
 - Sampling of 40 trees / treatment with deadwood
- Landscape scale analysis
 - Buffer around sampling units (1, 2.5 and 5 km)



PARTIAL RESULTS

Saproxylic beetles




Figure 2. Mean number of saproxylic beetles as a function of treatments and year. Different letters refer to significant differences ($\alpha = 0.05$). Asterisk indicates that all means of 2016 are different than those of 2015. Method: Linear mixed-effects model.




Figure 3. Total number of saproxylic beetle (Log) as a function of the proportion of old-growth/perturbed forest within a 5 km buffer around sampling units. Asterisk indicates significant slopes ($\alpha = 0.05$). Method: Linear mixed-effects model.

Feeding marks

Table 1. Type III tests of fixed effects on occurrence probability of woodpecker feeding marks ($\alpha = 0.05$). Method: General linear model (Logit link).

Fixed effects	Estimate	SD	P
Treatment	4.693	1.633	0.004
Proportion of old-growth/perturbed forest	0.250	0.067	<0.001
Mean DBH of black spruce in vegetation plot	0.766	0.319	0.016
Total number of saproxylic beetles at the nearest trap	0.001	0.0004	0.005
Forest stand density	-0.001	0.001	0.324
DHP of the sampled tree	0.001	0.042	0.987
Natural deadwood on vegetation plot	-0.028	0.032	0.383
Natural deadwood on vegetation plot *Treatment	-0.213	0.071	0.003




Figure 4. Interaction between the probability of woodpecker feeding mark presence ($\pm 95\%$ CI) and the proportion of old-growth/perturbed forest within a 5 km buffer around sampling units. Method: General linear model (Logit link).

Nest boxes

- 6 Boreal Chickadee Successes
- Two smaller nest boxes

Point counts

- Diversity index (2015 vs. 2016)
- Multivariate analyses




Figure 5. Example of a fresh woodpecker feeding mark on a girdled black spruce

References: ¹ Östlund, L., Zackrisson, O., and Axelsson, A.-L. 1997. Canadian Journal of Forest Research. 27: 1198-1206. ² Boucher, J., Azeria, E.T., Ibarzabal, J., and Hébert, C. 2012. Ecoscience. 19(4): 328-343. ³ Cyr, D., Gauthier, S., Bergeron, Y., and Carcaillat, C. 2009. Frontiers in Ecology and the Environment. 7(10): 519-524. ⁴ Fridman, J., and Walheim, M. 2000. Forest Ecology and Management. 131: 23-36. ⁵ Roberge, J.M., Lămás, T., Lundmark, T., Ranius, T., Felton, A., and Nordin, A. 2015. Journal of Environmental Management. 154: 284-292. ⁶ Vaillancourt, M.A., Drapeau, P., Gauthier, S., and Robert, M. 2008. Forest Ecology and Management. 255(7): 2272-2285. ⁷ Imbeau, L., Mönkkönen, M., and Desrochers, A. 2001. Conservation Biology. 15(4): 1151-1162. ⁸ Drapeau, P., Leduc, A., Bergeron, Y., Gauthier, S., and Savard, J.P. 2003. Forestry Chronicle. 79(3): 531-540. ⁹ Crête, M., Brais, S., Campagna, M., Darveau, M., Despons, M., Déry, S., Drapeau, P., Drolet, B., Jetté, J.-P., Maisonneuve, C., Nappi, A., and Petitclerc, P. 2004. Ministère des Ressources Naturelles, 35 p. ¹⁰ Kilgo, J.C., and Vukovich, M.A. 2014. Biological Conservation. 171: 21-28. ¹¹ Seibold, S., Bässler, C., Brandl, R., Gossner, M.M., Thorn, S., Ulyshen, M.D., and Müller, J. 2015. Biological Conservation. 191: 139-149. ¹² Lambrechts and al. 2010. Acta Ornithologica. 45(1): 1-26. ¹³ Aitken, K.E.H., and Martin, K. 2012. The Journal of Wildlife Management. 76(3): 557-565. ¹⁴ Robles, H., Ciudad, C., and Matthysen, E. 2012. Forest Ecology and Management. 277: 46-53. ¹⁵ Underwood, A.J. 1994. Ecological Applications. 4(1): 3-15.